SECTION E

INDEX TO VOLUME 22

Authors

Boyd, E. M.—Species variation in normal blood lipids estimated by oxidative micromethods, 39.

Cameron, A. T.—The relative sweetnesses of certain sugars, mixtures of sugars, and glycerol, 45.

Cleghorn, R. A. and McKelvey, A. D.—Studies of shock produced by muscle trauma. I. Methods; mortality; cardiovascular, blood concentration, and sugar changes, 12.

Collier, H. B. and Mack, G. E., Jr.—Vitamin B and phenothiazine anaemia in dogs, 1.

Craigie, J.—An alkaline thionin method of staining rickettsiae, 89.

Dowding, E. S.—The spores of dermatophytes, 67.

Ettinger, G. H.—See Sawyer, M. E. M.

Mack, G. E. Jr.—See Collier, H. B.

McKelvey, A. D.—See Cleghorn, R. A.

Melville, K. I. and Stehle, R. L.—Chemotherapy in experimental tuberculosis, 95.

Noble, R. L.—See Toby, C. G.

Reynolds, A. K. and Waud, R. A.—The pharmacological actions of capaurine, 64.

Sawyer, M. E. M. and Ettinger, G. H.—The cardiac action of posterior pituitary extract in physiological doses, in the normal dog, and after partial and complete denervation of the heart, 26. *(This paper is a reprint of one published in Can. J. Research, D, 21: 311-323. 1943.)*

Stehle, R. L.—See Melville, K. I.

Toby, C. G. and Noble, R. L.—Factors affecting the development of resistance to traumatic shock in the rat, 79.

Waud, R. A.—See Reynolds, A. K.

SECTION E

INDEX TO VOLUME 22

Subjects

Alcohol, See Glycerol.

Aleuriospores, See under Spores.

Alkaloid(s), See Capaurine.

Anaemia, Phenothiazine, in dogs, Vitamin B as related to, 1.

Bacteria, See Tuberculosis.

Blood
- concentration changes accompanying production of shock by muscle trauma, 17.
- lipids, Variations in, in different species of animals, as estimated by oxidative micromethods, 39.
- pressure
  - of dog
    - as affected by posterior pituitary extract, 26.
    - as affected by shock produced by muscle trauma, 16.
  - of rabbit, Effect of capaurine on, 65.
- sugar changes accompanying production of shock by muscle trauma, 20.

Bullfrog, Blood lipids of, as estimated by oxidative micromethods, 39.

Capaurine and its O-methyl ether derivative, Pharmacological action of, 64.

Cardiac action
- of capaurine, 65.
- of posterior pituitary extract in physiological doses, in the normal dog, and after partial and complete denervation of the heart, 26.

Cardiovascular changes
- accompanying production of shock by muscle trauma, 16.
- See also Cardiac action.

Chemotherapy
- in experimental tuberculosis, 95.
- Vitamin B as related to phenothiazine anaemia in dogs, 1.

Chlamydospores, See under Spores.

Circulation of frog and rabbit, as affected by capaurine, 65.

Conidia, See under Spores.

Corydalis aurea, Alkaloid from, and its pharmacological action, 64.

Dermatophytes, See under Fungi.

Dog(s)
- Blood lipids of, as estimated by oxidative micromethods, 39.
- Normal and after partial and complete denervation of the heart, Cardiac action of posterior pituitary extract in, 26.
- Studies of shock produced by muscle trauma in, 12.
- Vitamin B and phenothiazine anaemia in, 1.

Epidermophyton, See under Fungi.

Frogs, Pharmacological action of capaurine on, 64.

Fumariaceae, Alkaloid from a member of, and its pharmacological action, 64.

Fungi
- Dermatophytes
  - Species of, in Canada, 69.
  - Spores of, as criteria for differentiation of species, 67.
- Epidermophyton floccosum
  - Only species in Canada, 69.
  - Spore morphology of, 73.
- Microsporum
  - Species of, in Canada, 69.
  - Spore morphology as basis of differentiation of species of, as compared to other criteria, 71, 73.
- Trichophyton
  - Species of, in Canada, 69.
  - Spore morphology as basis of differentiation of species of, as compared to other criteria, 71, 73.

Glucose, Relative sweetness of, when compared with other sugars, mixtures of sugars, and glycerol, 45.

Glycerol, Sweetness of, as compared to certain sugars and mixtures of sugars, 45.

Guinea pigs, Tests of the effect of 79 chemicals on tuberculosis induced in, 99.

Haemoconcentration, See Blood concentration.

Heart
of dog
after bilateral vagotomy, Effect of posterior pituitary extract on, 34.
after complete denervation, Effect of posterior pituitary extract on, 36.
after thoracic sympathectomy, Effect of posterior pituitary extract on, 34.
after unilateral vagotomy and bilateral sympathectomy, Effect of posterior pituitary extract on, 35.
as affected by shock produced by muscle trauma, 16.
normal, Effect of posterior pituitary extract on, 28.
of frog, Effect of capaurine on, 65.

Heart rate, See Cardiovascular changes.

Lactose, Relative sweetness of, when compared with other sugars, mixtures of sugars, and glycerol, 45.

Lipids in blood of different species of animals, Variations in, 39.

Macroconidia, See under Spores.

Mice, Pharmacological action of capaurine on, 64.

Microconidia, See under Spores.

Micromethods, Oxidative, Use of, in estimation of blood lipids, 39.

Microsporum, See under Fungi.

Micro-organisms, See Rickettsiae and Tuberculosis.

Mortality
in dogs, accompanying production of shock by muscle trauma, 14.
in rats, as affected by traumatic shock, 82.

Muscles of certain animals, Action of capaurine on, 66.

Muscle trauma, See Shock.

Oidia, See under Spores.

Pharmacological action of capaurine, 64.

Phenothiazine anaemia in dogs, Vitamin B as related to, 1.

Pituitary extract, Cardiac action of, in the normal dog and after partial and complete denervation of the heart, 26.

Rabbit(s), Pharmacological action of capaurine on, 65.

Rana catesbiana, See Bullfrog.

Rat, Factors affecting resistance to traumatic shock in, 79.

Resistance to traumatic shock in the rat, Factors affecting, 79.

Rickettsiae, An alkaline thionin method of staining, 89.

Shock
Production of, by muscle trauma
Blood concentration changes accompanying, 17.
Blood sugar changes accompanying, 20.
Cardiovascular changes accompanying, 16.
Methods used in study of, 13.
Mortality accompanying, 14.
Traumatic
in the rat, Factors affecting development of resistance to, 79.
See also Shock, Production of.

Species variation in normal blood lipids estimated by oxidative micromethods, 39.

Spirochaete(s), Possible use of an alkaline thionin stain on, 92.

Spores of dermatophytes
as criteria for differentiation of species, 67.
Definition of kinds of
Aleuriospores, 67.
Chlamydospores, 67.
Conidia, 67.
Macroconidia, 67.
Microconidia, 67.
Oidia, 67.

Stain, Alkaline thionin, for rickettsiae, 89.

Sucrose, Relative sweetness of, when compared with other sugars, mixtures of sugars and glycerol, 45.

Sugar(s)
Blood, See Blood sugar.
Sweetness of, as compared to other sugars, mixtures of sugars, and glycerol, 45.

Sweetness, Relative, of certain sugars, mixtures of sugars, and glycerol, 45.

Thionin method, Alkaline, of staining rickettsiae, 89.

Trauma
in the rat, Factors affecting resistance to shock caused by, 79.
Muscle, See Shock.

Trichophyton, See under Fungi.

Tubercle bacillus, See Tuberculosis.

Tuberculosis induced experimentally in guinea pigs, Tests of the effect of 79 chemicals on, 99.

Virus(es), Possible use of an alkaline thionin stain on, 92.

Vitamin B as related to phenothiazine anaemia in dogs, 1.